

(b) a fragment of SEQ ID NO:2 consisting of at least amino acids 120 to 127 of SEQ ID NO:2.

71. (Three Times Amended) An isolated polynucleotide molecule comprising a nucleic acid 95% or more identical to a reference nucleic acid encoding an amino acid sequence selected from the group consisting of:

- (a) amino acids 1 to 127 of SEQ ID NO:2;
- (b) amino acids 2 to 127 of SEQ ID NO:2; and
- (c) the amino acid sequence encoded by the cDNA clone contained in ATCC

Deposit No. 97856;

wherein said polynucleotide is overexpressed in breast cancer.

72. (Twice Amended) An isolated polynucleotide comprising a nucleic acid encoding an amino acid sequence, wherein, except for one to thirty conservative amino acid substitutions, said amino acid sequence is selected from the group consisting of:

- (a) amino acids 1 to 127 of SEQ ID NO:2;
- (b) amino acids 2 to 127 of SEQ ID NO:2; and
- (c) the amino acid sequence encoded by the cDNA clone contained in ATCC

Deposit No. 97856;

wherein said polynucleotide is overexpressed in breast cancer.

76. (Twice Amended) An isolated polynucleotide comprising a nucleic acid encoding an amino acid sequence 95% or more identical to a reference amino acid sequence from the group consisting of:

- (a) amino acids 1 to 127 of SEQ ID NO:2;
- (b) amino acids 2 to 127 of SEQ ID NO:2; and
- (c) the amino acid sequence encoded by the cDNA clone contained in ATCC

Deposit No. 97856;

wherein said polynucleotide is overexpressed in breast cancer.

77. (Three Times Amended) An isolated polynucleotide comprising a nucleic acid which is 95% or more identical to a reference nucleic acid, wherein said reference nucleic acid is selected from the group consisting of:

- (a) nucleotides 15 to 392 of SEQ ID NO:1; and

(b) nucleotides 12 to 392 of SEQ ID NO:1;
wherein said polynucleotide is overexpressed in breast cancer.
